

Infrared-suppressed gluon propagator in 4d Yang-Mills theory in a Landau-like gauge

Attilio Cucchieri^a Axel Maas^{a,b} Tereza Mendes^a

^a*Instituto de Física de São Carlos, Universidade de São Paulo, Caixa Postal 369,
13560-970 São Carlos, SP, Brazil*

^b*Gesellschaft für Schwerionenforschung mbH, Planckstraße 1, D-64291
Darmstadt, Germany*

Abstract

The infrared behavior of the gluon propagator is directly related to confinement in QCD. Indeed, the Gribov-Zwanziger scenario of confinement predicts an infrared vanishing (transverse) gluon propagator in Landau-like gauges, implying violation of reflection positivity and gluon confinement. Finite-volume effects make it very difficult to observe (in the minimal Landau gauge) an infrared suppressed gluon propagator in lattice simulations of the four-dimensional case. Here we report results for the $SU(2)$ gluon propagator in a gauge that interpolates between the minimal Landau gauge (for gauge parameter λ equal to 1) and the minimal Coulomb gauge (corresponding to $\lambda = 0$). For small values of λ we find that the spatially-transverse gluon propagator $D^{\text{tr}}(0, |\vec{p}|)$, considered as a function of the spatial momenta $|\vec{p}|$, is clearly infrared suppressed. This result is in agreement with the Gribov-Zwanziger scenario and with previous numerical results in the minimal Coulomb gauge. We also discuss the nature of the limit $\lambda \rightarrow 0$ (complete Coulomb gauge) and its relation to the standard Coulomb gauge ($\lambda = 0$). Our findings are corroborated by similar results in the three-dimensional case, where the infrared suppression is observed for all considered values of λ .

Key words: Yang-Mills theory, Gluon propagator, Confinement, Interpolating gauge

PACS: 11.15.Ha, 12.38.Aw, 14.70.Dj

Email addresses: attilio@ifsc.usp.br (Attilio Cucchieri), axelmaas@web.de (Axel Maas), mendes@ifsc.usp.br (Tereza Mendes).

1 Introduction

The infrared behavior of the gluon propagator is linked to the confinement of gluons.[1] In particular, the confinement scenario of Gribov and Zwanziger[2,3,4,5,6,7,8] predicts a (transverse) gluon propagator vanishing at zero (Euclidean) momentum in Landau gauge and in Landau-like gauges (or λ -gauges). The latter class refers to gauges interpolating between the Landau and the Coulomb gauge[9], with a gauge condition (in d dimensions) given by

$$\lambda \partial_0 A_0^a + \partial_1 A_1^a + \dots + \partial_{d-1} A_{d-1}^a = 0, \quad (1)$$

where the gauge parameter λ is between 1 and 0. Let us recall that an infrared (IR) null gluon propagator has far-reaching consequences. Indeed, such a particle cannot have a positive semi-definite spectral function[10,11,12,13] or, as a consequence, a Källen-Lehmann representation. This is regarded as one possible manifestation of confinement,[1,14,15] when considering Euclidean correlation functions.¹

The question of whether the Landau-gauge gluon propagator is indeed null at zero momentum is a long-standing one. Various continuum methods, based on functional approaches, yield a vanishing gluon propagator.[1,6,7,8,16,17,18,19,20,21,22,23,24,25] This result is rather tightly constrained,[25] i.e. it seems to be the only possible solution satisfying both Dyson-Schwinger equations and functional renormalization-group equations. At the same time, lattice calculations in four dimensions have obtained an IR-suppressed Landau-gauge gluon propagator $D(p)$ only when using strongly-asymmetric lattices[26] or a coupling constant in the strong-coupling regime.[27] For lattice couplings in the scaling region and using symmetric lattices, one finds for the Landau-gauge gluon propagator an increase slower than $1/p^2$ as one approaches the IR region[28,29,30,31], with a finite value for $p = 0$. [32,33,34,35] The fact that a propagator decreasing at small momenta is not observed in the 4d case, even for volumes of almost $(10 \text{ fm})^4$, [35] is probably due to very strong finite-size effects.[32,33,34,35,36,37,38,39] This assumption is supported by numerical results in the 3d case, where much larger lattice sides are accessible. In this case, there is substantial evidence for an IR-suppressed gluon propagator in Landau gauge,[40,41,42] in agreement with continuum calculations.[6,43] However, also in this case, a reliable extrapolation of $D(0)$ to the infinite-volume limit is still lacking.[42] Finally, let us recall that lattice Landau calculations[44,45,36,37] have also obtained direct evidence for the non-positivity of the gluon spectral function, both in the three- and in the four-dimensional cases.

The Gribov-Zwanziger confinement scenario applies also to Coulomb gauge.[2,3,4,46,47,48,49,50] In this case it is important to observe that the standard Coulomb-gauge-fixing

¹ Thus, it is a sufficient condition for gluon confinement.[1,15]

condition $\partial_1 A_1^a + \dots + \partial_{d-1} A_{d-1}^a = 0$ is not a complete one, due to the residual gauge degrees of freedom $g(t)$. On the other hand, a possible complete Coulomb gauge condition[9] can be obtained using the class of gauges defined in (1). Indeed, the parameter λ interpolates between the Landau ($\lambda = 1$) and a complete Coulomb gauge, corresponding to the limit² $\lambda \rightarrow 0$. Therefore, the complete Coulomb gauge condition is, by definition, a smooth limiting case of the interpolating gauge (1) while, of course, this is not the case for the standard Coulomb condition ($\lambda = 0$). Let us recall that the gauge condition (1) above can be obtained by minimizing the (lattice) functional³

$$\mathcal{E}[g] = -\text{Tr} \sum_x \left[\lambda U_0(x) + \sum_{i=1}^{d-1} U_i(x) \right], \quad (2)$$

where $U_\mu(x)$ indicates a lattice link variable in the μ direction. Then, the limiting case $\lambda \rightarrow 0$ corresponds to minimizing the following two functionals[51]

$$\mathcal{E}_{\text{hor}}[g(\vec{x})] = -\text{Tr} \sum_x \sum_{i=1}^{d-1} U_i(x) \quad (3)$$

$$\mathcal{E}_{\text{ver}}[g(t)] = -\text{Tr} \sum_x U_0(x). \quad (4)$$

The minimization of the first functional is equivalent to a Landau gauge condition fixed on each time slice, using $g(\vec{x})$ gauge transformations, i.e. it corresponds to the standard (incomplete) Coulomb gauge. The minimization of the second functional, considering only $g(t)$ gauge transformations, provides additional constraints, necessary to eliminate the residual gauge degrees of freedom. Note that we can also write $\mathcal{E}_{\text{ver}}[g(t)] = -\text{Tr} \sum_t Q_0(t)$, with $Q_0(t) = \sum_{\vec{x}} U_0(t, \vec{x})$. Then, the minimization of $\mathcal{E}_{\text{ver}}[g(t)]$ is like a one-dimensional Landau gauge fixing. Of course, quantities defined in terms of spatial link variables $U_i(x)$ are not affected by the residual gauge condition obtained by minimizing the functional $\mathcal{E}_{\text{ver}}[g(t)]$.

Numerical studies in minimal Coulomb gauge have shown [for the $SU(2)$ case in 4d] that the instantaneous transverse gluon propagator $D^{tr}(\vec{p})$ is indeed suppressed in the IR limit.[51,52,53,54,55] Also, in the infinite-volume limit, it has been found[51,52,53] that $D^{tr}(\vec{p})$ is well described by a Gribov-like propagator with a pair of purely imaginary poles $m^2 = \pm i\gamma$. These results are in agreement with the Gribov-Zwanziger confinement scenario.[2,3,4,46,47,48] The fact that, for a given lattice size L (in fm), one sees an IR-suppressed transverse gluon propagator in 4d-Coulomb gauge and in 3d-Landau gauge but not in 4d-Landau gauge may be related to a quantitatively different IR

² Clearly, since the gauge fixing (1) is complete for any $\lambda \neq 0$, it is also a complete one when considering the limit $\lambda \rightarrow 0$. [9]

³ A similar functional can be defined in the continuum.

suppression in the two cases. Indeed, functional methods[6,43,56,57,58,59] predict a stronger suppression in 4d-Coulomb gauge and in 3d-Landau gauge than in 4d-Landau gauge, i.e. the so-called IR gluon exponent α_D should be larger for the 4d-Coulomb and the 3d-Landau cases.

One should note that, for any non-zero value of λ , the gauge condition (1) is essentially a *deformed* Landau gauge, i.e. the IR exponents of the propagators do not depend on λ [60]. In particular, calculations using Dyson-Schwinger equations suggest[60] that, at momenta sufficiently small compared to a separation momenta p_s , the transverse gluon propagator behaves as in Landau gauge, i.e. all Lorentz and color components of the gluon propagator vanish at zero four-momentum. However, the limit $\lambda \rightarrow 0$ can also be thought of as sending to zero the momentum p_s , which separates Coulomb-gauge-like from Landau-gauge-like behavior. Indeed, for any finite value $\lambda \neq 0$ and given the Landau gauge condition $\partial_\mu A_\mu^a = 0$, we can obtain the gauge condition

$$\partial_0 A_0^a + \lambda^{-1} \left[\partial_1 A_1^a + \dots + \partial_{d-1} A_{d-1}^a \right] = 0 \quad (5)$$

by using the rescaling⁴ $x_i \rightarrow x_i/\lambda$ for $i \neq 0$. This implies, in momentum space, the rescaling $p_i \rightarrow \lambda p_i$. Thus, if we consider only spatial momenta, we find⁵ that p_s is rescaled to λp_s and goes to zero when $\lambda \rightarrow 0$. As a consequence, one should expect that, for very small values of λ , all correlation functions would show a Coulomb-like behavior for momenta $p > p_s$, with p_s very small. In particular, the correlation function that corresponds to the transverse (instantaneous) gluon propagator in Coulomb gauge should become more and more IR suppressed as the parameter λ approaches zero. Investigating whether this is the case is the aim of this work.

2 Numerical results

Following Eqs. (8) and (9) of Ref. [60], we can consider on the lattice the three-dimensionally transverse gluonic correlation function

$$D^{\text{tr}}(p_0, |\vec{p}|) = \left(\delta_{ij} - \frac{p_i p_j}{\vec{p}^2} \right) \frac{\langle A_i^a(p) A_j^a(-p) \rangle}{(N_c^2 - 1)(d - 2)V} . \quad (6)$$

At zero four-momentum one has

$$D^{\text{tr}}(0) = \delta_{ij} \frac{\langle A_i^a(0) A_j^a(0) \rangle}{(N_c^2 - 1)(d - 1)V} . \quad (7)$$

⁴ In Ref. [60], a similar rescaling was used to show that Dyson-Schwinger equations for λ -gauges are equivalent to the Landau case for all $\lambda \neq 0$.

⁵ Of course, this simple explanation is correct only at tree-level and can be modified by the renormalization of λ .

Here, N_c is the number of colors, V is the d -dimensional lattice volume, Lorentz indices i, j are summed only over the $d - 1$ spatial directions and $A_i^a(p)$ is the d -dimensional Fourier transform of the gluon field. Let us recall that, by considering only spatial momenta (i.e. $p_0 = 0$), the function $D^{\text{tr}}(0, |\vec{p}|)$ is predicted to be IR suppressed — and vanishing at zero momentum — for all non-zero values of λ in three and in four dimensions.[60] Also, when λ is null, the above definition yields the instantaneous part of the three-dimensional transverse gluon propagator[51]

$$D^{\text{inst}}(|\vec{p}|) = \sum_t \left(\delta_{ij} - \frac{p_i p_j}{\vec{p}^2} \right) \frac{\langle A_i^a(t, \vec{p}) A_j^a(t, -\vec{p}) \rangle}{(N_c^2 - 1)(d - 2)V}, \quad (8)$$

where now the Fourier transform of the gluon field is evaluated for each time slice. Indeed, when λ is null, the gauge transformations $g(t)$ are independent of the gauge transformation $g(\vec{x})$ and the two sets of transformations commute. Then, the terms in Eqs. (6) that depend explicitly on $g(t)$ are averaged to zero and one is left with the expression above. Note that this explanation is valid whether the residual gauge freedom $g(t)$ is fixed or not.

Here we evaluate numerically $D^{\text{tr}}(0, |\vec{p}|)$ as a function of $|\vec{p}|$ for $SU(2)$ Yang-Mills theory in three and in four dimensions for several values of the parameter λ . Details of the simulations can be found in.[42,61] Let us note that for $\lambda \neq 1$ the numerical gauge fixing is very similar to the usual Landau gauge fixing.[41] On the other hand, when λ goes to zero one sees[62] that more iterations are needed in order to satisfy a given numerical accuracy for the gauge fixing, in agreement with a recent analytic study.[63] This problem can be partially reduced by adding some extra gauge fixing sweeps in the $\mu = 0$ direction for each iteration of the gauge-fixing algorithm, i.e. by considering gauge transformations $g(t)$ that depend only on the $\mu = 0$ component of x . Finally, we did not consider here possible systematic effects related to the breaking of rotational symmetry or to the existence of Gribov copies. The former type of effects can be parameterized by[64] $a^2 p^{[4]}$, where a is the lattice spacing and $p^{[4]} = \sum_\mu p_\mu^4$. Therefore, this type of effects are not expected to play a significant role in the IR region, considered here. As for the latter type of effects, in the infinite-volume limit, averages taken over configurations belonging to the so-called Gribov region Ω should coincide[8] with averages obtained by restricting the functional integral to the so-called fundamental modular region Γ , whose interior is free of Gribov copies.

Our results⁶ are reported in Figure 1. In three dimensions, a well-defined maximum (and thus an IR suppression) is visible for all values of λ . This includes also Landau gauge ($\lambda = 1$), confirming earlier results.[40,41,42] As can be seen from the plot, the maximum value attained by the propagator seems to move to larger momenta with decreasing λ , going from about 400

⁶ Preliminary results have been presented in Ref. [65,66].




Fig. 1. The gluonic correlation function $D^{\text{tr}}(0, |\vec{p}|)$ in three (top figure) and four (bottom figure) dimensions. The spatial momenta $|\vec{p}|$ are chosen along a single axis. Circles indicate data for $\lambda = 1$ (Landau gauge), crosses are used for $\lambda = 1/2$, squares for $\lambda = 1/10$, triangles for $\lambda = 1/20$, stars for $\lambda = 1/100$ and upside-down triangles represent results at $\lambda = 0$ (Coulomb gauge). Data have been obtained at $\beta = 4.2$ in three dimensions and at $\beta = 2.2$ in four dimensions. The lattice size is $40^3 \approx (6.9 \text{ fm})^3$ in three dimensions and $22^4 \approx (4.6 \text{ fm})^4$ in four dimensions. In addition, in the bottom panel, diamonds correspond to a $40^4 \approx (8.4 \text{ fm})^4$ lattice at $\lambda = 1/100$. The physical scale has been set using Refs. [42,67]. Note that our error bars are smaller than the sizes of the symbols. Also, for $\lambda = 0$, we consider the standard (incomplete) Coulomb gauge.




Fig. 2. Same as in Fig. 1, except that here we show data for the $D_{00}(0, |\vec{p}|)$ tensor component.

MeV in the Landau case[42] to about 600 MeV for the Coulomb case. At the same time, as λ decreases, the maximum becomes more visible. In four dimensions, on the other hand, no discernible peak is visible in Landau gauge when considering the volume $V = 22^4$. Decreasing λ , however, leads to a suppression of the propagator in the IR region. In particular, at the smallest value of λ considered, i.e. $\lambda = 1/100$ and for a lattice volume $V = 40^4$, a maximum is seen also in four dimensions. This maximum is as visible as the one in three dimensions in Landau gauge. Thus for small λ , just as for Coulomb gauge,[51] one sees a maximum of the transverse gluon propagator already for

relatively small volumes.

A similar λ -dependence is also seen in the tensor component $D_{00}(0, |\vec{p}|)$. In particular, as λ becomes small, this tensor component of the gluon propagator becomes enhanced for small (nonzero) values of $|\vec{p}|$ (see Fig. 2). Let us recall that, in Coulomb gauge, the D_{00} component is IR divergent already at the perturbative level. Moreover, the instantaneous part of D_{00} is related to the color Coulomb potential and should be diverging as $1/|\vec{p}|^4$ at small momenta.[46,47,48,51,52,53,54,55] Thus, from the discussion presented in the Introduction, one could expect to observe a D_{00} component enhanced at intermediate momenta $p > p_s$ but still suppressed at small momenta. This is indeed the case and from the plots in Fig. 2 one clearly sees how this enhancement is developing for $|\vec{p}| \neq 0$ as the value of λ decreases. Let us stress that the discontinuity in the behavior of the propagator at $|\vec{p}| \neq 0$ is observed since we are taking the limit $\lambda \rightarrow 0$. A similar discontinuity is also obtained when implementing the complete Coulomb gauge described in the Introduction.⁷ On the other hand, for the incomplete Coulomb gauge, D_{00} is clearly a continuous function of $|\vec{p}|$ (see Fig. 2).

As said in the Introduction, the IR exponent α_D for the transverse correlator is predicted to show a discontinuity in the limit $\lambda \rightarrow 0$. This is not seen from our data. However, a reliable check of this prediction can only be obtained if one has control over the infinite-volume and the continuum limits, which we have not yet achieved. On the other hand, taking the limit $\lambda \rightarrow 0$ clearly induces a discontinuity in the behavior of the temporal component D_{00} . [60]

3 Conclusions

We have presented the first direct observation of an IR-suppressed gluonic correlation function on a symmetric 4d lattice in a Landau-like gauge, with gauge parameter λ as in Eq. (1). (Landau gauge is obtained for $\lambda = 1$, while Coulomb gauge corresponds to $\lambda = 0$.) The suppression is seen for sufficiently small λ when considering moderately small lattice volumes. Judging from the results shown for the 3d case, it is conceivable that a similar suppression might be observed for any λ if a large enough lattice side is considered. Furthermore, since the limit $\lambda \rightarrow 1$ is smooth,[9,60] we expect to see an IR suppression for sufficiently large lattices also in Landau gauge. Moreover, we have obtained that the D_{00} component of the gluon propagator gets enhanced in the IR region (for $|\vec{p}| \neq 0$) at small values of the interpolating parameter λ . These results clearly demonstrate the transformation of the correlation functions from Landau gauge towards Coulomb gauge when considering the limit $\lambda \rightarrow 0$. They

⁷ This is a consequence of the minimization of $\mathcal{E}_{\text{ver}}[g(t)]$, defined in Eq. (4).

also provide additional support to the Gribov-Zwanziger scenario of confinement, establishing an IR suppression of gluonic correlation functions in four dimensions beyond Coulomb gauge. The use of the interpolating gauge thus constitutes a promising alternative to studies in Landau gauge, since finite-size effects are significantly smaller, allowing a more efficient investigation of the gauge dependence of correlation functions of confined objects.

Acknowledgments

A. M. was supported in part by the DFG under grant number MA 3935/1-1. A. C. and T. M. were supported by FAPESP (under grants # 00/05047-5 and 05/59919-7) and by CNPq. The ROOT framework[68] has been used in this project. Part of our simulations has been done on the IBM supercomputer at São Paulo University (FAPESP grant # 04/08928-3).

References

- [1] R. Alkofer and L. von Smekal, Phys. Rept. **353** (2001) 281 [arXiv:hep-ph/0007355].
- [2] V. N. Gribov, Nucl. Phys. B **139** (1978) 1.
- [3] D. Zwanziger, Phys. Lett. B **257** (1991) 168.
- [4] D. Zwanziger, Nucl. Phys. B **364** (1991) 127.
- [5] D. Zwanziger, Nucl. Phys. B **412** (1994) 657.
- [6] D. Zwanziger, Phys. Rev. D **65** (2002) 094039 [arXiv: hep-th/0109224].
- [7] D. Zwanziger, Phys. Rev. D **67** (2003) 105001 [arXiv: hep-th/0206053].
- [8] D. Zwanziger, Phys. Rev. D **69** (2004) 016002 [arXiv: hep-ph/0303028].
- [9] L. Baulieu and D. Zwanziger, Nucl. Phys. B **548** (1999) 527 [arXiv:hep-th/9807024].
- [10] R. Oehme and W. Zimmermann, Phys. Rev. D **21** (1980) 471.
- [11] R. Oehme and W. Zimmermann, Phys. Rev. D **21** (1980) 1661.
- [12] K. Nishijima, Int. J. Mod. Phys. A **9** (1994) 3799.
- [13] K. Nishijima, Int. J. Mod. Phys. A **10** (1995) 3155.
- [14] J. E. Mandula, Phys. Rept. **315** (1999) 273 [arXiv:hep-lat/9907020].

- [15] R. Alkofer, W. Detmold, C. S. Fischer and P. Maris, Phys. Rev. D **70** (2004) 014014 [arXiv:hep-ph/0309077].
- [16] L. von Smekal, A. Hauck and R. Alkofer, Phys. Rev. Lett. **79** (1997) 3591 [arXiv:hep-ph/9705242].
- [17] C. Lerche and L. von Smekal, Phys. Rev. D **65**, 125006 (2002) [arXiv:hep-ph/0202194].
- [18] L. von Smekal, A. Hauck and R. Alkofer, Annals Phys. **267** (1998) 1 [Erratum-ibid. **269** (1998) 182] [arXiv:hep-ph/9707327].
- [19] C. S. Fischer and R. Alkofer, Phys. Lett. B **536** (2002) 177 [arXiv:hep-ph/0202202].
- [20] C. S. Fischer, R. Alkofer and H. Reinhardt, Phys. Rev. D **65** (2002) 094008 [arXiv:hep-ph/0202195].
- [21] H. Gies, Phys. Rev. D **66** (2002) 025006 [arXiv:hep-th/0202207].
- [22] C. S. Fischer and H. Gies, JHEP **0410** (2004) 048 [arXiv:hep-ph/0408089].
- [23] J. M. Pawłowski, D. F. Litim, S. Nedelko and L. von Smekal, Phys. Rev. Lett. **93** (2004) 152002 [arXiv:hep-th/0312324].
- [24] R. Alkofer, C. S. Fischer and F. J. Llanes-Estrada, Phys. Lett. B **611** (2005) 279 [arXiv:hep-th/0412330].
- [25] C. S. Fischer and J. M. Pawłowski, Phys. Rev. D **75** (2007) 025012 [arXiv:hep-th/0609009].
- [26] P. J. Silva and O. Oliveira, Phys. Rev. D **74** (2006) 034513 [arXiv:hep-lat/0511043].
- [27] A. Cucchieri, Phys. Lett. B **422** (1998) 233 [arXiv:hep-lat/9709015].
- [28] F. Gutbrod, DESY-96-252;
- [29] A. Cucchieri, Nucl. Phys. B **508** (1997) 353 [arXiv:hep-lat/9705005].
- [30] D. B. Leinweber, J. I. Skullerud, A. G. Williams and C. Parrinello, Phys. Rev. D **58** (1998) 031501(R) [arXiv:hep-lat/9803015].
- [31] S. Furui and H. Nakajima, Phys. Rev. D **69** (2004) 074505 [arXiv:hep-lat/0305010].
- [32] J. C. R. Bloch, A. Cucchieri, K. Langfeld and T. Mendes, Nucl. Phys. B **687** (2004) 76 [arXiv:hep-lat/0312036].
- [33] A. Sternbeck, E. M. Ilgenfritz, M. Müller-Preussker and A. Schiller, Phys. Rev. D **72** (2005) 014507 [arXiv:hep-lat/0506007].
- [34] P. Boucaud *et al.*, arXiv:hep-lat/0602006.
- [35] A. Cucchieri and T. Mendes, arXiv:hep-ph/0605224.

- [36] A. Sternbeck, arXiv:hep-lat/0609016, Ph. D. Thesis (Humboldt Universität, Berlin).
- [37] A. Sternbeck, E. M. Ilgenfritz, M. Muller-Preussker, A. Schiller and I. L. Bogolubsky, PoS **LAT2006** (2006) 076 [arXiv:hep-lat/0610053].
- [38] C. S. Fischer, A. Maas, J. M. Pawłowski and L. von Smekal, arXiv:hep-ph/0701050.
- [39] A. Cucchieri, A. Maas and T. Mendes, Phys. Rev. D **75**, 076003 (2007) [arXiv:hep-lat/0702022].
- [40] A. Cucchieri, Phys. Rev. D **60** (1998) 034508 [arXiv:hep-lat/9902023].
- [41] A. Cucchieri, F. Karsch and P. Petreczky, Phys. Rev. D **64** (2001) 036001 [arXiv:hep-lat/0103009].
- [42] A. Cucchieri, T. Mendes and A. R. Taurines, Phys. Rev. D **67** (2003) 091502(R) [arXiv:hep-lat/0302022].
- [43] A. Maas, J. Wambach, B. Grüter and R. Alkofer, Eur. Phys. J. C **37**, No.3 (2004) 335 [arXiv:hep-ph/0408074].
- [44] K. Langfeld, H. Reinhardt and J. Gattnar, Nucl. Phys. B **621** (2002) 131 [arXiv:hep-ph/0107141].
- [45] A. Cucchieri, T. Mendes and A. R. Taurines, Phys. Rev. D **71** (2005) 051902(R) [arXiv:hep-lat/0406020].
- [46] D. Zwanziger, Nucl. Phys. B **518** (1998) 237.
- [47] A. Cucchieri and D. Zwanziger, Phys. Rev. D **65** (2002) 014002.
- [48] D. Zwanziger, Phys. Rev. Lett. **90** (2003) 102001.
- [49] Y. L. Dokshitzer and D. E. Kharzeev, Ann. Rev. Nucl. Part. Sci. **54** (2004) 487.
- [50] A. Cucchieri, AIP Conf. Proc. **892** (2007) 22 [arXiv:hep-lat/0612004].
- [51] A. Cucchieri and D. Zwanziger, Phys. Rev. D **65** (2002) 014001.
- [52] A. Cucchieri and D. Zwanziger, Phys. Lett. B **524** (2002) 123.
- [53] A. Cucchieri, T. Mendes and D. Zwanziger, Nucl. Phys. Proc. Suppl. **106** (2002) 697.
- [54] A. Cucchieri and D. Zwanziger, Nucl. Phys. Proc. Suppl. **119** (2003) 727.
- [55] K. Langfeld and L. Moyaerts, Phys. Rev. D **70** (2004) 074507.
- [56] D. Zwanziger, Phys. Rev. D **70** (2004) 094034 [arXiv: hep-ph/0312254].
- [57] C. Feuchter and H. Reinhardt, Phys. Rev. D **70** (2004) 105021 [arXiv:hep-th/0408236].

- [58] H. Reinhardt and C. Feuchter, Phys. Rev. D **71** (2005) 105002 [arXiv:hep-th/0408237].
- [59] W. Schleifenbaum, M. Leder and H. Reinhardt, Phys. Rev. D **73** (2006) 125019 [arXiv:hep-th/0605115].
- [60] C. S. Fischer and D. Zwanziger, Phys. Rev. D **72** (2005) 054005 [arXiv:hep-ph/0504244].
- [61] A. Cucchieri, A. Maas and T. Mendes, Phys. Rev. D **74** (2006) 014503 [arXiv:hep-lat/0605011].
- [62] A. Cucchieri and T. Mendes, arXiv:hep-lat/9902024.
- [63] A. Cucchieri and T. Mendes, Comput. Phys. Commun. **154** (2003) 1 [arXiv:hep-lat/0301019].
- [64] J. P. Ma, Mod. Phys. Lett. A **15** (2000) 229 [arXiv:hep-lat/9903009].
- [65] A. Maas, A. Cucchieri and T. Mendes, arXiv:hep-lat/0610123.
- [66] Braz. J. Phys. **37N1B** (2007) 219 [arXiv:hep-lat/0610006].
- [67] J. Fingberg, U. M. Heller and F. Karsch, Nucl. Phys. B **392** (1993) 493 [arXiv:hep-lat/9208012].
- [68] R. Brun and F. Rademakers, Nucl. Instrum. Meth. A **389** (1997) 81.